

ACTA BIOMEDICA REPRINT

ATENEI PARMENSIS | FOUNDED 1887

Official Journal of the Society of Medicine and Natural Sciences of Parma

*The Acta Biomedica is indexed by Index Medicus / Medline Excerpta Medica (EMBASE),
the Elsevier BioBASE*

POSITIVE CLINICAL OUTCOMES DERIVED FROM USING A PROPRIETARY MIXTURE OF SELECTED STRAINS DURING PREGNANCY

Francesco Di Pierro, Annamaria Parolari, Benedetta Brundu, Raffaele Nigro



MATTIOLI 1885



Positive clinical outcomes derived from using a proprietary mixture of selected strains during pregnancy

Francesco Di Pierro¹, Annamaria Parolari², Benedetta Brundu³, Raffaele Nigro²

¹Scientific Department, Velleja Research, Milan, Italy; ²O.U. Gynecology and Obstetrics Department, Arco Hospital, Trento, Italy; ³O.U. Gynecology and Obstetrics, Maternal-fetal Medicine, "G. Saliceto" Hospital, Piacenza, Italy

Summary. *Background:* While the use of antibiotic prophylaxis is clearly advantageous to prevent streptococcal infection, it alters the composition of the gut microbiota in mothers and infants. *Enterococcus faecium* L3 is one of the best studied probiotic strains and shows strong antagonistic activity against *Streptococcus agalactiae* due to the production of bacteriocins able to inhibit common gut and vaginal pathogens. *Methods:* We tested an L3-based probiotic formula (iNatal®) on 127 pregnant women attending our gynaecological unit in 2015. We compared the study subjects with 279 pregnant women enrolled in the same year and with 892 other pregnant women who attended our gynaecological unit in 2013 and 2014. *Results:* The findings demonstrate: (a) the safety profile of the product; (b) its ability to reduce gut disorders; (c) a 6% decrease in the incidence of streptococcal colonization; (d) an approximately 30% decline in episodes of premature rupture of membranes; (e) fewer caesarean sections during labour; and (f) a reduction in pathological umbilical cord blood pH. *Conclusions:* Our results demonstrate that a probiotic treatment during pregnancy could have unexpected but favourable clinical results. Further randomized, double-blind, placebo controlled studies are now needed to confirm our preliminary findings. (www.actabiomedica.it)

Key words: *Enterococcus faecium* L3, PROM, *Streptococcus agalactiae*, Intrapartum Antibiotic Prophylaxis, iNatal

Introduction

Streptococcus agalactiae, a gram-positive commensal bacterium present in the gastrointestinal and genitourinary tracts of 10-30% of pregnant women, is the primary source of early-onset bacterial sepsis in newborns, a major cause of neonatal morbidity and mortality (1). The newborn can be exposed to streptococcus in utero, as the organism can reach the amniotic fluid through ruptured membranes and/or the birth canal during delivery. The introduction in many countries of universal screening for the presence of streptococcus in all pregnant women and subsequent intrapartum ampicillin prophylaxis in *S. agalactiae*-positive women has

resulted in a significant reduction in early-onset infection, which has fallen from 1.7 to 0.25 cases per 1000 births in developed countries (2). While the use of antibiotic prophylaxis is clearly advantageous for the prevention of streptococcal infection, it alters the composition of the gut microbiota of infants and results in a higher relative abundance of ampicillin-resistant *Enterobacteriaceae* and in a dramatic reduction in bifidobacteria (3). Recently, it has been suggested that mother-only supplementation with probiotics during pregnancy could reduce vaginal carriage of *S. agalactiae* (4). One of the best studied probiotic strains with strong activity against *S. agalactiae* is *Enterococcus faecium* L3. Its antagonistic activity is due to the produc-

tion of two well-known bacteriocins called enterocin A and enterocin B (5). In addition to *S. agalactiae*, this bacteriocin-releasing strain has also shown activity against the *in vitro* growth of *Escherichia coli*, *Salmonella* spp., *Shigella* spp., *Staphylococcus aureus*, *Proteus vulgaris*, *Streptococcus pyogenes*, *Klebsiella pneumonia*, *Mycoplasma hominis*, *Candida albicans* and *Helicobacter pylori* (6). Strain L3 is also pharmacologically active *in vitro* against herpes simplex virus type 1 (HSV-1). In particular, light and immunofluorescence microscopy showed an anti-virus effect on HSV-1 grown on Vero cell cultures in the presence of strain L3 or in the presence of its two bacteriocins (7). Clinically, in mature newborns receiving antibiotic therapy (penicillins, aminoglycosides, cephalosporin), administration of *E. faecium* L3 reduced the incidence and persistence of *Clostridium difficile* infection as well as a number of opportunistic microorganisms in the intestinal microbiota while preserving the growth of bifidobacteria and lactobacilli (8). In premature infants receiving full or partial parenteral nutrition and antimicrobial therapy (carbapenems, fluoroquinolones, glycopeptides), the use of strain L3 resulted in increased body weight, decreased the frequency of infectious complications, promoted conservation of immunomodulatory function and restricted the growth of nosocomial flora (9,10). A recent paper has also shown that the use of *E. faecium* L3 in children in their first year of life reduced the incidence of acute respiratory infections in the winter-spring period by more than 50% and improved the infants' nutritional status resulting in a significantly higher body mass index (11). Finally, a pilot uncontrolled study performed on 112 pregnant women indicated that strain L3 had anti-candida and anti-streptococcal properties. Administered during the last trimester, the strain reduced gynaecological infections (12). In this study, we examined whether the antibiotic-like properties of *E. faecium* L3, clearly reported as *in vitro* results or seen clinically in mature and preterm newborns, would also be demonstrated if administered to women during pregnancy. We have therefore evaluated whether administration of a probiotic mixture containing strain L3 during the last 10 weeks of pregnancy could affect outcomes such as the occurrence of *S. agalactiae* or premature rupture of membranes (PROM).

Materials and Methods

Methods

The trial was conducted at the Gynecology and Obstetrics Department, Arco Hospital, Trento, Italy and in accordance with the Declaration of Helsinki, with the approval of the local ethics committee. Each participant was told of the study procedures and objectives and signed an informed consent form and a form covering personal data handling in accordance with privacy law. The objectives of the study were to examine: (a) the safety and tolerability of a probiotic treatment; (b) the number of rectal-vaginal swabs positive for *S. agalactiae*; (c) episodes of PROM; (d) the rate of caesarean sections during labour; and (e) umbilical cord arterial blood pH.

Inclusion and exclusion criteria

The study enrolled 406 healthy pregnant women attending the department from January to December 2015. Exclusion criteria included: age below 18; the presence of a neurological disorder, heart, lung or kidney disease, or a severe metabolic disorder; a past history of cancer; and refusal to sign the informed consent and/or privacy form.

Group assignment

We performed a non-randomized, controlled, open-label study. The probiotic product, iNatal®, was administered to 127 pregnant women, while 279 served as controls. Subjects were assigned to the treatment or control group as follows. Of the 406 women attending our department during 2015, 166 reported having episodes of constipation, colitis, diarrhoea or bladder infections in the trimester before enrolment or had evidence of vaginosis, vulvo-vaginitis or undiagnosed vulvar itching. These 166 women were offered treatment with the probiotic product. Of these 166 women, 127 accepted and were assigned to the treated group, while the remaining 39 and the other 240 women not reporting any gut or gynaecological disorder in the previous trimester served as controls (untreated).

Treatment

iNatal® was administered daily 2 hours after breakfast from the 30th to the 40th week of pregnancy. Acetaminophen and antibiotics could be taken if required. However, the probiotic treatment was suspended during antibiotic therapy but was then resumed and continued until the scheduled end day. iNatal® is a nutritional supplement notified at the Italian Ministry of Health by Omeopiacenza (Pontenure, Italy), according to the provisions of law No. 169 of 2004, on 4 September 2014 (notification number: 71640) and contains *Enterococcus faecium* L3 LMG P-27496 at the concentration of 5×10^9 colony forming units/dose (CFU/dose), *Bifidobacterium animalis* ssp. *lactis* BB12 DSM 15954 at 3×10^9 CFU/dose, *Lactobacillus casei* RO215 CNCM I-3429 at 3×10^9 CFU/dose and *Lactococcus lactis* ssp. *lactis* SP 38 DSM 26868 at 3×10^9 CFU/dose.

Swab test

A rectal-vaginal swab (Todd Hewitt CNA Regular Flocked Swab Kit; Copan Diagnostics, Murrieta, CA, USA) was taken between the 36th and 37th weeks of pregnancy to check for the presence of *S. agalactiae*.

Statistical analysis

Equivalence between the two groups (treated and control) regarding age, parity, ethnicity, education and type of employment was determined using Fisher's exact test and the two-tailed Wilcoxon-Mann-Whitney test. The difference between the two groups in terms of side effects and clinical outcomes was determined using the two-tailed Wilcoxon-Mann-Whitney test. JMP 10 statistical software for Mac OS X and statistical significance was set at 95%.

Results

A total of 406 pregnant women were enrolled in a non-randomized, controlled, open label clinical study during which 127 pregnant women were administered a proprietary mixture of selected probiotic strains, while

279 served as controls. Despite the large difference in numbers of subjects in the two groups, statistical analysis showed that the groups were comparable as regards age, parity, ethnicity, education and employment (Table 1). The study was carried out to examine whether the administration of iNatal® was safe and well tolerated, and could reduce the number of rectal-vaginal swabs positive for *S. agalactiae*, the occurrence of PROM, the rate of caesarean sections during labour and the presence of pathological umbilical cord arterial blood pH. As shown in Table 2, the product was found to be safe and well tolerated with no particular side effects. It is noteworthy that gut side effects, such as bowel pain, constipation, colitis, meteorism, flatulence and diarrhoea, were less frequent in the treated group than in control subjects. This finding is likely due to the probiotic effect of the tested formula. Rectal-vaginal swabs were positive for the presence of streptococcus in 27 of

Table 1. Characteristics of the 406 women enrolled in the study and attending our unit in 2015

	Treated (127)	Untreated (279)	p Value*
Age			
18-19	0	1	ns
20-24	12	31	ns
25-29	30	70	ns
30-34	37	83	ns
35-39	34	69	ns
40-44	14	25	ns
Children			
1	56	125	ns
2	37	99	ns
3	34	55	ns
Ethnicity			
European	82	186	ns
North African	28	62	ns
Far Eastern	11	20	ns
Other	6	11	ns
Education			
Primary school	1	2	ns
Middle school	34	59	ns
High school	65	139	ns
University	27	79	ns
Employment			
None/housewife	43	97	ns
Yes	84	182	ns

*Non-significant (ns) differences between groups

Table 2. Adverse events (number of global episodes) during 10-week treatment with the proprietary mixture of selected strains and in the control group

	Gastric ¹	Gut ²	Skin ³	Headache	Insomnia
Treated (127)	60	11*	1	15	68
Control (279)	136	75	3	39	152

¹Pain, reflux, nausea, vomiting, spasm; ²pain, constipation, colitis, meteorism, flatulence, diarrhoea; ³rash, erythema, dermatitis, acne-like reaction.

*p<0.01 versus control.

the 127 women in the treated group (21.3%) and in 76 of the 279 (27.3%) women in the control group (Table 3). The significant reduction in swab positivity is confirmed when positive findings rates in the treated 2015 group are compared with the higher rates in untreated women from two previous years (2014 and 2013). No episodes of PROM (Table 4) were seen in the treated group compared to 87 episodes in the control group (31.2%), which findings were similar to the 156/290 (40%) and 180/502 (35.9%) PROM episodes in the previous two years. In 2015, seven (5.51%) and 29 (10.39%) caesarean sections were performed during labour in the treated and control groups, respectively, because of concerning cardiotocography (CTG) findings (Table 5). According to the Robson classification (13) the seven caesarean sections of the treated group were ranked as Group 1 as well as 25 out of 29 of the control one. The remaining four of the control group were ranked as Group 3 (data not shown). The number of caesarean sections in the untreated group is very similar to the 43/390 cases (11.02%) in 2014 and 57/502 cases (11.35%) in 2013. Finally, we considered pathological the umbilical cord arterial blood pH when lower than 7.2. As shown in Table 6, there were no cases of blood pH <7.2 in the treated group but there were 17

Table 3. Number of women attending our gynaecological department in 2013-2015 with rectal-vaginal swabs positive for *Streptococcus agalactiae*

	Treated 2015	Untreated 2015	All 2014	All 2013
Positive/total %	27/127* 21.3	76/279 27.3	125/390 32.05	138/502 27.5

*p<0.05 versus Untreated 2015, All 2014 and All 2013.

Table 4. Episodes of PROM^o in all women attending our gynaecological department, 2013-2015

	Treated 2015	Untreated 2015	All 2014	All 2013
PROM/total %	0/127* 0.0	87/279 31.2	156/390 40	180/502 35.9

*p<0.001 versus Untreated 2015, All 2014 and All 2013.

^oPROM: premature rupture of membranes.

Table 5. Caesarean sections during labour in all women attending our gynaecological department, 2013-2015

	Treated 2015	Untreated 2015	All 2014	All 2013
Caesarean/total %	7/127* 5.51	29/279 10.39	43/390 11.02	59/502 11.75

*p<0.05 versus Untreated 2015, All 2014 and All 2013.

Table 6. Umbilical cord arterial blood pH <7.2 in all women attending our gynaecological department in 2013-2015

	Treated 2015	Untreated 2015	All 2014	All 2013
<7.2/total %	0/127* 0.0	17/279 6.09	20/390 5.12	29/502 5.79*

p<0.01 versus Untreated 2015, All 2014 and All 2013.

cases out of 279 (6.09%) in the control group. Values in the two previous years were comparable to those in the control group, with 20/390 (5.12%) in 2014 and 29/502 (5.79%) in 2013.

Discussion

The use of intrapartum antibiotics likely affects the biodiversity of neonatal microbiota and is associated with mother and infant gut microbiota dysbiosis (14). Alteration of gut microbiota in infants caused by caesarean section can have later negative consequences such as chronic immune disease, allergic rhinitis, atopy, asthma, overweight and obesity (15-18). Since intrapartum antibiotics are often administered because of rectal and/or vaginal colonization by *S. agalactiae*, we have tested the role of *E. faecium* L3, a probiotic select-

ed strain known to counteract streptococcal *in vitro* growth, in reducing the clinical occurrence of *S. agalactiae* as evaluated by rectal-vaginal swab. No pharmaceutical or nutritional supplement-grade products containing just *E. faecium* L3 are available in Italy as the strain is only offered in a probiotic product also containing *Bifidobacterium animalis* ssp. *lactis* BB12, *Lactobacillus casei* RO215 and *Lactococcus lactis* ssp. *lactis* SP 38. Strain BB12 has been intensively investigated. It has immune-modulating properties (19), is capable of potentiating a specific IgG response to vaccination (20) and can reduce the incidence of upper respiratory tract infections (21). *L. casei* is a probiotic described as having anti-inflammatory properties likely mediated by IL-10 (22) and useful for potentiating the role of mesalazine in diverticular disease (23). In particular, strain RO215 (data not shown) is a strong producer (29.8 mg/L) of L-tryptophan, which is converted to melatonin in the gastrointestinal tract during the day when pineal gland synthesis is inhibited (24). Clinically, gastrointestinal melatonin has potential for the prevention or treatment of ulcerative colitis, irritable bowel syndrome, childhood colic diarrhoea and necrotizing enterocolitis in premature infants (25,26). *L. lactis* ssp. *lactis*, previously known as *Streptococcus lactis* (27), has lactose-fermenting ability and proteinase activity likely mediated through two distinct plasmids (28). On the basis of these findings, the L3-based mixture of strains likely confers benefits on both pregnant women and their infants. Probiotic intervention during pregnancy is an opportunity to promote the health of both the mother and the child (29). Probiotic strains administered to pregnant woman can be found in the infant gut and temporary colonization of an infant may be possible by colonizing the pregnant mother before delivery. Colonization is stable for as long as 6 months, and in unexplained circumstances may persist for up to 24 months (30). Administration of specific probiotics to pregnant women positive for the skin prick test was found to be a safe and effective method for reducing the risk of eczema in infants evaluated at 6, 12 and 24 months (31). Our study has shown that administration of iNatal® is safe and well tolerated. As expected, the mixture reduces the gut discomfort typically affecting pregnant women, and therefore acts as a eubiotic. The mixture also decreases swab positivity for *S. agalacti-*

ae, thereby reducing the use of intrapartum antibiotics. We expected a greater decrease, but some recent findings could explain why only a 6% reduction was observed (32,33). In fact, the very high colonization index (100% in healthy female volunteers; data not shown) of the enterococcus strain used in the product may be partially responsible for the minimal result being the enterococcus strain confused with *S. agalactiae*. However, following iNatal® treatment, only 21.3% of women in the treated group required intrapartum antibiotics compared to 27.3% in the control group. Episodes of PROM were also reduced in the treated group. PROM and preterm premature rupture of membranes (pPROM) are associated with an elevated risk of intra-amniotic infection which increases perinatal morbidity and mortality. Nevertheless, PROM is only very rarely life-threatening and it might be caused by gynaecological dysbiotic conditions (34). We believe that the reduction in the number of PROM episodes is mainly due to the antimicrobial effects of the L3 strain which is a strong producer of bacteriocins, although the immuno-modulating properties of BB12, or some other beneficial and still unknown actions of the other strains, cannot be excluded. Finally, we do not know why use of the probiotic product reduced the number of unplanned caesarean sections during labour or why the treatment reduced the incidence of umbilical cord blood pH <7.2. However, we suggest that the product improved the general condition of the mother and fetus. Our work has of course a number of limitations. First, the two groups are very different in terms of numbers and subjects are not randomized. This difference, and the lack of randomization, are anyway due to the established criteria for enrolling the women to be treated. As clearly shown in Table 1, in spite of this difference the two groups are then absolutely comparable according to their main features. Second, our study is not double-blind and placebo controlled. In spite of these two last important bias, our study shows an important number of controls. The outcomes measured have been in fact compared with 1171 untreated pregnant women attending our operative unit from 2013 to 2015. Nevertheless, our results prompt us to perform a new clinical trial with randomization, blinding and the presence of a probiotic treatment for the control group.

Conflict of interest

FDP is a member of the scientific council of a company selling iNatal®. The other authors have no conflicts of interest.

References

- Johri AK, Paoletti LC, Glaser P, et al. Group B Streptococcus: global incidence and vaccine development. *Nat Rev Microbiol* 2006; 4: 932-42.
- Rivera L, Sáez-Llorens X, Feris-Iglesias J, et al. Incidence and serotype distribution of invasive group B streptococcal disease in young infants: a multi-country observational study. *BMC Pediatr* 2015; 15: 143-49.
- Mazzola G, Murphy K, Ross RP, et al. Early gut microbiota perturbations following intrapartum antibiotic prophylaxis to prevent Group B streptococcal disease. *PLoS One* 2016; 11(6): e0157527.
- Barthow C, Wickens K, Stanley T, et al. The Probiotics in Pregnancy Study (PiP Study): rationale and design of a double-blind randomised controlled trial to improve maternal health during pregnancy and prevent infant eczema and allergy. *BMC Pregnancy Childbirth* 2016; 16(1): 133.
- Ermolenko EI, Chernysh AI, Martsinkovskaia IV, Suvorov AN. Influence of probiotic enterococci on the growth of *Streptococcus agalactiae*. *Zh Mikrobiol Epidemiol Immunobiol* 2007; (5): 73-7.
- Di Pierro F. Bacterial therapy in pregnancy to drive the intra-partum colonization of newborns. *Nutrafoods* 2106; 15: 5-8.
- Ermolenko EI, Furaeva VA, Isakov VA, Ermolenko DK, Suvorov AN. Inhibition of herpes simplex virus type 1 reproduction by probiotic bacteria in vitro. *Vopr Virusol* 2010; 55(4): 25-8.
- Lo Skiavo LA, Gonchar NV, Fedorova MS, Suvorov AN. Dynamics of contamination and persistence of *Clostridium difficile* in intestinal microbiota in newborn infants during antibiotic therapy and use of probiotic strain *Enterococcus faecium* L3. *Antibiot Khimioter* 2013; 58(11-12): 13-18.
- [No authors listed]. Value of probiotic in reducing frequency of infectious complications in premature infants. *Antibiot Khimioter* 2014; 59(1-2): 30-5.
- Gonchar NV, Alekhina LA, Suvorov AN. Probiotic strains of enterococci as a means of therapy and prophylaxis of intestinal diseases in children (review of literature). *Eksp Klin Gastroenterol* 2013; (1): 74-8.
- Gonchar NV, Suvorov AN, Maryshev VP, Sorokina TM, Churkova TV, Kharit SM. Probiotics, nutritional status and resistance to respiratory infections in infants. *Eksp Klin Gastroenterol* 2015; (1): 48-54.
- Di Pierro F. Probiotici specifici in gravidanza per pilotare la colonizzazione intra-partum del nascituro [Specific probiotics for pregnancy to pilot intra-partum colonization of newborn]. *Int Nut* 2014; 17(4): 2-6.
- Robson MS. Classification of caesarean sections. *Fetal and Maternal Medicine Review* 2001; 12(1): 23-39.
- Azad MB, Konya T, Persaud RR, et al.; CHILD Study Investigators. Impact of maternal intrapartum antibiotics, method of birth and breastfeeding on gut microbiota during the first year of life: a prospective cohort study. *BJOG* 2016; 123(6): 983-93.
- Sevelsted A, Stokholm J, Bønnelykke K, Bisgaard H. Cesarean section and chronic immune disorders. *Pediatrics* 2015; 135(1): e92-8.
- Pistiner M, Gold DR, Abdulkarim H, Hoffman E, Celedón JC. Birth by cesarean section, allergic rhinitis, and allergic sensitization among children with a parental history of atopy. *J Allergy Clin Immunol* 2008; 122(2): 274-9.
- Thavagnanam S, Fleming J, Bromley A, Shields MD, Cardwell CR. A meta-analysis of the association between Caesarean section and childhood asthma. *Clin Exp Allergy* 2008; 38(4): 629-33.
- Huh SY, Rifas-Shiman SL, Zera CA, Edwards JW, Oken E, Weiss ST, et al. Delivery by caesarean section and risk of obesity in preschool age children: a prospective cohort study. *Arch Dis Child* 2012; 97(7): 610-16.
- Holscher HD, Czerkies LA, Cekola P, et al. *Bifidobacterium lactis* Bb12 enhances intestinal antibody response in formula-fed infants: a randomized, double-blind, controlled trial. *JPEN J Parenter Enteral Nutr* 2012; 36 (1 Suppl): 106S-17S.
- Rizzardini G, Eskesen D, Calder PC, Capetti A, Jespersen L, Clerici M. Evaluation of the immune benefits of two probiotic strains *Bifidobacterium animalis* ssp. *lactis*, BB-12® and *Lactobacillus paracasei* ssp. *paracasei*, *L. casei* 431® in an influenza vaccination model: a randomised, double-blind, placebo-controlled study. *Br J Nutr* 2012; 107(6): 876-84.
- Taipale TJ, Pienihäkkinen K, Isolauri E, Jokela JT, Söderling EM. *Bifidobacterium animalis* subsp. *lactis* BB-12 in reducing the risk of infections in early childhood. *Pediatr Res* 2016; 79(1-1): 65-9.
- Qiu ZB, Chen J, Chen JJ, et al. Effect of recombinant *Lactobacillus casei* expressing interleukin-10 in dextran sulfate sodium-induced colitis mice. *J Dig Dis* 2013; 14(2): 76-83.
- Tursi A, Brandimarte G, Elisei W, et al. Randomized clinical trial: mesalazine and/or probiotics in maintaining remission of symptomatic uncomplicated diverticular disease--a double-blind, randomized, placebo-controlled study. *Aliment Pharmacol Ther* 2013; 38(7): 741-51.
- Zagajewski J, Drozdowicz D, Brzozowska I, et al. Conversion L-tryptophan to melatonin in the gastrointestinal tract: the new high performance liquid chromatography method enabling simultaneous determination of six metabolites of L-tryptophan by native fluorescence and UV-VIS detection. *J Physiol Pharmacol* 2012; 63(6): 613-21.
- Bubenik GA. Localization, physiological significance and possible clinical implication of gastrointestinal melatonin. *Biol Signals Recept* 2001; 10(6): 350-66.
- Marseglia L, D'Angelo G, Manti S, et al. Oxidative stress-mediated damage in newborns with necrotizing entero-

- colitis: a possible role of melatonin. *Am J Perinatol* 2015; 32(10): 905-9.
27. Chopin MC, Chopin A, Rouault A, Galleron N. Insertion and amplification of foreign genes in the *Lactococcus lactis* subsp. *lactis* chromosome. *Appl Environ Microbiol* 1989; 55(7): 1769-74.
28. Efstathiou JD, McKay LL. Plasmids in *Streptococcus lactis*: evidence that lactose metabolism and proteinase activity are plasmid linked. *Appl Environ Microbiol* 1976; 32(1): 38-44.
29. Isolauri E, Rautava S, Collado MC, Salminen S. Role of probiotics in reducing the risk of gestational diabetes. *Diabetes Obes Metab* 2015; 17(8): 713-19.
30. Schultz M, Göttl C, Young RJ, Iwen P, Vanderhoof JA. Administration of oral probiotic bacteria to pregnant women causes temporary infantile colonization. *J Pediatr Gastroenterol Nutr* 2004; 38(3): 293-7.
31. Rautava S, Kainonen E, Salminen S, Isolauri E. Maternal probiotic supplementation during pregnancy and breastfeeding reduces the risk of eczema in the infant. *J Allergy Clin Immunol* 2012; 130(6): 1355-60.
32. Savini V, Gherardi G, Marrollo R, et al. Could β -hemolytic, group B *Enterococcus faecalis* be mistaken for *Streptococcus agalactiae*? *Diagn Microbiol Infect Dis* 2015; 82(1): 32-3.
33. Savini V, Franco A, Gherardi G, et al. Beta-hemolytic, multi-lancefield antigen-agglutinating *Enterococcus durans* from a pregnant woman, mimicking *Streptococcus agalactiae*. *J Clin Microbiol* 2014; 52(6): 2181-2.
34. Veleminský M, Tosner J. Relationship of vaginal microflora to PROM, pPROM and the risk of early-onset neonatal sepsis. *Neuro Endocrinol Lett* 2008; 29(2): 205-21.
-
- Received: 19 July 2016
Accepted: 25 August 2016
Correspondence:
Francesco Di Pierro,
Velleja Research, Viale Lunigiana 23,
20125, Milan, Italy
Tel. +39 3495527663
Fax: + 39 0523 511894.
E-mail: f.dipierro@vellejaresearch.com

ACTA BIO MEDICA

ATENEI PARMENSIS

FOUNDED 1887

OFFICIAL JOURNAL OF THE SOCIETY OF MEDICINE AND NATURAL SCIENCES OF PARMA

free on-line: www.actabiomedica.it

EDITOR IN CHIEF

Maurizio Vanelli - Parma, Italy

DEPUTY EDITORS

Loris Borghi - Parma, Italy

Marco Vitale - Parma, Italy

HONORARY EDITORS

Carlo Chezzi - Parma, Italy

Roberto Delsignore - Parma, Italy

Almerico Novarini - Parma, Italy

Giacomo Rizzolatti - Parma, Italy

EDITORIAL BOARD

Fernando Arevalo - Caracas, Venezuela
Judy Aschner - Nashville, TN, USA
Michael Aschner - Nashville, TN, USA
Franco Aversa - Parma, Italy
Alberto Bacchi Modena - Parma, Italy
Cesare Beghi - Varese, Italy
Corrado Betterle - Padova, Italy
Mauro Bonanini - Parma, Italy
Loris Borghi - Parma, Italy
David A. Bushinsky - Rochester, NY, USA
Ovidio Bussolati - Parma, Italy
Ardeville Cabassi - Parma, Italy
Paolo Caffarra - Parma, Italy
Anthony Capone Jr. - Detroit, MI, USA
Francesco Ceccarelli - Parma, Italy
Gian Paolo Ceda - Parma, Italy
Graziano Ceresini - Parma, Italy
Gianfranco Cervellin - Parma, Italy
Alfredo Antonio Chetta - Parma, Italy
Marco Colonna - St. Louis, MO, USA
Paolo Coruzzi - Parma, Italy
Lucio Guido Maria Costa - Parma, Italy
Cosimo Costantino - Parma, Italy
Giuseppe Crovini - Fidenza, Italy
Domenico Cucinotta - Bologna, Italy
Alessandro De Fanti - Reggio Emilia, Italy
Massimo De Filippo - Parma, Italy
Filippo De Luca - Messina, Italy

Giuseppe Fabrizi - Parma, Italy
Valentina Fainardi - Parma, Italy
Guido Fanelli - Parma, Italy
Claudio Feliciani - Parma, Italy
Antonio Freyrie - Parma, Italy
Vittorio Gallese - Parma, Italy
Livio Garattini - Milano, Italy
Mario J. Garcia - New York, NY, USA
Donald J. Hagler - Rochester, MINN, USA
Rick Hippakka - Chicago, IL, USA
Andrew R. Hoffman - Stanford, CA, USA
Joachim Klosterkoetter - Colonia, Germany
Ingrid Kreissig - Heidelberg, Germany
Ronald M. Lechan - Boston, MA, USA
Annarosa Leri - Harvard, Boston, MA, USA
Giuseppe Lippi - Verona, Italy
Nicola Longo - Salt Lake City, UT, USA
Wanyun Ma - Beijing, China
Umberto Vittorio Maestroni - Parma, Italy
Marcello Giuseppe Maggio - Parma, Italy
Norman Maitland - York, United Kingdom
Federico Marchesi - Parma, Italy
Carla Mastrorilli - Parma, Italy
James A. McCubrey - Greenville, NC, USA
Tiziana Meschi - Parma, Italy
Antonio Mutti - Parma, Italy
Giuseppe Nuzzi - Parma, Italy
Jose Luis Navia - Cleveland, OH, USA

Donald Orlic - Bethesda, MD, USA
Marc S. Penn - Cleveland, OH, USA
Silvia Pizzi - Parma, Italy
Francesco Pogliacomini - Parma, Italy
Federico Quaini - Parma, Italy
Edoardo Raposio - Parma, Italy
Stephen M. Rao - Cleveland, OH, USA
Luigi Roncoroni - Parma, Italy
Shaukat Sadikot - Mumbai, India
Simone Cherchi Sanna - New York, NY, USA
Leopoldo Sarli - Parma, Italy
Robert S. Schwartz - Denver, Colorado, USA
Anthony Seaton - Edinburgh,
United Kingdom
Mario Sianesi - Parma, Italy
Carlo Signorelli - Parma, Italy
Mario Strazzabosco - New Haven, CT, USA
Nicola Sverzellati - Parma, Italy
Maria Luisa Tanzi - Parma, Italy
Roberto Toni - Parma, Italy
Frederik H. Van Der Veen - Maastricht,
The Netherlands
Vincenzo Vincenti - Parma, Italy
Vincenzo Violi - Parma, Italy
Richard Wallensten - Solna, Sweden
Francesco Ziglioli - Reggio Emilia, Italy

LINGUISTIC ADVISOR

Rossana Di Marzio
Parma, Italy

EDITORIAL OFFICE MANAGER

Anna Scotti
Mattioli 1885 srl - Casa Editrice
Strada di Lodesana 649/sx, Loc. Vaio
43036 Fidenza (PR), Italy
Tel. ++39 0524 530383
Fax ++39 0524 82537
contact@actabiomedica.it

Francesco Covino
Società di Medicina e Scienze Naturali
Azienda Ospedaliero-Universitaria
di Parma - Cattani Building, 2nd floor
Via Gramsci, 14 - Parma, Italy
Tel./Fax ++39 0521 033730
francesco.covino@unipr.it

PUBLISHER

Mattioli 1885 srl Casa Editrice
Strada di Lodesana, 649/sx, Loc. Vaio
43036 Fidenza (PR), Italy
Tel. ++39 0524 530383
Fax ++39 0524 82537
E-mail: edit@mattioli1885.com