

ORIGINAL ARTICLE

Use of a probiotic mixture containing *Bifidobacterium animalis* subsp. *lactis* BB12 and *Enterococcus faecium* L3 in atopic children

Francesco DI PIERRO ¹*, Isabella BASILE ², Maria L. DANZA ³,
Leo VENTURELLI ⁴, Raffaele CONTINI ⁵, Paolo RISSO ⁶, Maria COLOMBO ⁵

¹Scientific and Research Department, Velleja Research, Milan, Italy; ²ASL District 6, Cologno Monzese, Milan, Italy; ³ASL District 6, Binasco, Milan, Italy; ⁴ATS Bergamo, Bergamo, Italy; ⁵ATS Milano, Vignate, Milan, Italy; ⁶Department of Health Sciences, University of Genoa, Genoa, Italy

*Corresponding author: Francesco Di Pierro, Velleja Research, Viale Lunigiana 23, 20125, Milan, Italy.
E-mail: f.dipierro@vellejaresearch.com

ABSTRACT

BACKGROUND: Imbalance of the human gut microbiota in childhood, mainly due to low gut biodiversity and a low bifidobacterial load, has been suggested as a risk factor for atopy. Administration of *Enterococcus faecium* L3 in infants has been shown to increase the gut bifidobacterial count. The aim was to verify if a mixture of *Bifidobacterium animalis* subsp. *lactis* BB12 and *E. faecium* L3 could reduce the signs, symptoms and need for drugs in atopic children.

METHODS: We retrospectively analyzed, and compared with controls, clinical outcomes following use of BB12 and L3 strains when administered 3 months before or during the development of signs and symptoms of atopy.

RESULTS: When administered in the 3 months before the development of atopy, the BB12 and L3 strains significantly reduced ($P<0.001$) rhinitis, watery eyes and cough/bronchospasm. However, reduced efficacy was observed when the mixture was given during the 3 months of atopy. The mixture of strains also significantly reduced the use of oral antihistamines, inhaled corticosteroids (in the same children in two different years) and oral corticosteroids (in different children in the same year).

CONCLUSIONS: When administered as a prophylactic, the mixture of BB12 and L3 (iNatal Ped®) statistically decreases the signs and symptoms of atopy and reduces the use of drugs. Administration of the same probiotics as treatment after the appearance of atopy is less effective.

(Cite this article as: Di Pierro F, Basile I, Danza ML, Venturelli L, Contini R, Rizzo P, et al. Use of a probiotic mixture containing *Bifidobacterium animalis* subsp. *lactis* BB12 and *Enterococcus faecium* L3 in atopic children. Minerva Pediatrica 2018;70:418-24. DOI: 10.23736/S0026-4946.18.05203-9)

KEY WORDS: Hypersensitivity - Asthma - Rhinitis - Lacrimal apparatus diseases - Cough - Bronchial spasm.

The human gut microbiota consists of 500-1000 distinct bacterial species with millions of active genes capable of influencing the host immune system and the mechanisms of atopy.¹ Most of these bacterial species belong to five phyla: *Bacteroidetes*, *Firmicutes*, *Actinobacteria*, *Proteobacteria* and *Verrucomicrobia*, with *Bacteroidetes* and *Firmicutes* constituting more than 95% of all gut bacteria.² Imbalance of the

human gut microbiota in early childhood has been suggested as a risk factor for atopy.³ *Clostridium difficile* colonization early after birth, for instance, was associated with gasping during the first 6-7 years of life and with asthma at age 6-7 years.⁴ Moreover, children developing allergy in the first decade of life had less gut bacterial diversity in the first month after birth compared with non-allergic children,⁵ as well as a lower content

of *Bifidobacterium* and *Enterococcus* species.⁶ Lower relative abundance of bacteria such as *Bifidobacterium*, *Akkermansia* and *Faecalibacterium* in children has been implicated in childhood atopy and asthma⁷ and lower content in *Bifidobacterium* has been observed in some cases of adult asthma.⁸ A lot of evidence also supports the immunomodulatory effects of some bifidobacterial strains, such as *Bifidobacterium animalis* and *Bifidobacterium longum*. These strains seem to reduce the Th2 immune response and antagonize the pro-inflammatory LPS-driven cascades.⁹ *Enterococcus faecium* L3, a probiotic strain known for its ability to release the bacteriocins enterocin A and enterocin B which target gut and vaginal pathogens, when administered to newborn infants preserves and increases the growth of endogenous bifidobacterial species, in addition to reducing the number of opportunistic microorganisms.¹⁰ We therefore retrospectively evaluated whether the administration to atopic children of a mixture of the two well-documented probiotic strains *B. animalis* subsp. *lactis* BB12 and *E. faecium* L3 was able to reduce the signs and symptoms of atopy and/or to reduce the use of drugs administered to control asthma and allergy.

Materials and methods

Product

The probiotic mixture of *B. animalis* subsp. *lactis* BB12 (BB12) and *E. faecium* L3 (L3) was manufactured by Farmaceutici Procemsa (Nichelino, Turin, Italy) as single dose sachets. The product was notified to the Italian Ministry of Health on June 16, 2016 as iNatal Ped® by Omeopiaceenza (Pontenure, PC, Italy) according to the provisions of law No. 169 of 2004 (notification number: 87470). The preparation used in the study contained not less than 2 billion CFU/sachet of each strain (total dose: not less than 4 billion CFU/sachet) as declared in the expiry date.

Clinical trial

Our retrospective analysis was conducted on 89 (54 male and 35 female) children with atopy living in the Milan area of Italy (see Figure 1 for the study scheme). Some of the children (N.=46)




Figure 1.—Study scheme.

**Bifidobacterium animalis* subsp. *lactis* BB12 and *Enterococcus faecium* L3.

were treated with the probiotic mixture, while the remaining 43 served as untreated controls. In the prophylactic arm of the study, children were treated between January and March (2017) in the 3 months before the ‘allergic’ period (April–June, 2017), while in the treatment arm, children were treated during the allergic 3 months (April–June, 2017). As one of the researchers (MLD) joined the study in the middle of the prophylaxis 3 months, children enrolled by this physician (N.=22) were treated during the allergic 3 months. We compared the results in the children who received the mixture (N.=46) with the same children the year before (untreated 2016) and with another group of untreated children (N.=43) enrolled as a control group in 2017 (untreated 2017). During the allergic 3 months, all children (N.=89) were administered physician-prescribed conventional pediatric drugs to control atopy as required. The retrospective analysis was conducted in accordance with the Declaration of Helsinki and was approved by the local ethics committee in Milan (Italy). The parents of all study participants were informed of the study methods and signed the appropriate consent and privacy policy documents.

Inclusion and exclusion criteria

We included children living in the northern part of Milan, who were 2–14 years of age and attending one of our pediatric outpatient clinics with a diagnosis of atopy based on a skin prick test, radioallergosorbent (RAST) test and clinical examination. The children enrolled were all char-

acterized by atopy *versus* inhaled allergens occurring in the 3 months April-June. We excluded children who were not immunocompetent, had cardiomyopathy, intrinsic asthma or wheezing due to infective etiology, or current or previous use of desensitizing vaccines.

Treatments

The parents were asked not to give their children any other probiotics, including yogurt and dead bacterial cells (tyndallized bacteria, bacterial lysate, postbiotics) for the duration of the study. In case of antibiotic use, parents were told to stop probiotic administration and to resume it after the last antibiotic dose had been given. Acetaminophen or ibuprofen could be used to treat pain, in which case probiotic use could be continued. Parents were asked not to administer the probiotics when the child was hungry or had consumed a large meal so that the probiotics would survive better to colonize the gut. All children were treated with conventional antiatopic drugs during the 3 months April-June in 2017 according to their prescriptions (which were similar to those for the same 3 months in 2016). Any conditions or diseases compatible with the aim the study were managed in accordance the Italian pediatric guidelines. There were no cases of diseases incompatible with our study and all enrolled children completed the study.

Study aims

The primary endpoints of the study were tolerability, side effects and compliance with the probiotic treatment, and development of signs and symptoms of atopy (rhinitis, watery eyes, cough and bronchospasm). Symptoms were evaluated according to a visual analogue scale as follows: 0 = no signs; 1 = mild signs; 2 = moderate signs; 3 = severe signs. The secondary endpoint was the need for and use of conventional drugs such as oral and ophthalmic antihistamines, oral and inhaled corticosteroids, antileukotrienes and adrenergic β_2 agonists. Topical corticosteroids were not used since no child had a diagnosis of dermal atopy.

Statistical analysis

Aiming for a significance level of 10% and a power of 90%, with an effect size of at least 0.55, we determined that a sample size of at least 32 subjects was required in order that the 2017 prophylactic group could be compared with their 2016 untreated status. Aiming for a significance level of 10% and a power of 90%, with an effect size of 0.8, we determined that at least 60 subjects were required, 30 for each arm, in order that the 2017 treated and 2017 untreated group could be compared. Therefore, at least 60 subjects, 30 for each arm, needed to be enrolled.

The non-parametric Wilcoxon signed-rank test was used to compare the treated 2017 and untreated 2016 groups. The non-parametric Wilcoxon rank-sum test was used to compare the treated 2017 and untreated 2017 groups. Results are presented as the mean and standard deviation or the odds ratio, with 95% confidence intervals.

To determine the odds ratios, we created a new variable for the presence, or not, of symptoms or the use, or not, of conventional drugs during the observation period. Therefore, the variable did not indicate the number of times a symptom occurred or the number of days a drug was used. JMP 10 for Mac OS X was used for statistical analysis and statistical significance was set at 95%.

Results

Our aim was to determine if treatment with a mixture of *B. animalis* subsp. *lactis* BB12 and *E. faecium* L3 (BB12 and L3) in addition to being well tolerated, lacking side effects and having good compliance, was able to reduce the signs, symptoms and need for drug use in atopic children. As shown in Table I, II, children enrolled in the prophylactic and treatment arms of the study did not show any differences in age or sex. Similarly, there were no significant differences in birthweight, type of delivery (vaginal or cesarean), breastfeeding (or not), ethnic group (Caucasian or not), or attendance at a nursery, toy library or kindergarten (data not shown) between the groups. The BB12 and L3 mixture was very well tolerated by all children and did

TABLE I.—*Characteristics of children in the prophylaxis arm of the study.*

Group	Sex	N.	Age*	P
Untreated 2016	M	21	9.5±2.8	NS
	F	14	10.1±2.8	NS
Treated 2017	M	21	10±3.3	NS
	F	14	11.1±2.8	NS
Untreated 2107	M	18	8.1±2.3	NS
	F	14	8.4±3.4	NS

NS: not significant.

*Expressed in years (mean±standard deviation).

TABLE III.—*Signs and symptoms of atopy according to a visual analogue scale* in children in the prophylaxis arm of the study: comparison between treated and untreated children in the same year.*

Parameter	Untreated 2017	Treated 2017	Δ%	P
Rhinitis	2.3±1.1	1.2±1	-47.8	0.0001
Watery eyes	1.7±1.1	0.8±0.9	-52.9	0.0011
Cough/bronchospasm	2±1.1	1.1±1.1	-45.0	0.0019

*Expressed as mean±standard deviation with: 0 = no signs; 1 = mild signs; 2 = moderate signs; 3 = severe signs.

not cause any side effects. Compliance was also good and no child withdrew from the study (data not shown).

Children in the prophylactic arm showed a highly significant ($P<0.001$) reduction in signs and symptoms of atopy (Table III, IV). Rhinitis, watery eyes and cough/bronchospasm were reduced by approximately 50% in children treated in 2017 as compared with the same children untreated in 2016 and as compared with the other children untreated in the same 3 months of 2017.

In contrast, children in the treatment arm of the study only showed a tendency towards a reduction in signs and symptoms (Table V, VI). When different children treated or not in the

TABLE V.—*Signs and symptoms of atopy according to a visual analogue scale* in children in the treatment arm of the study: comparison between treated and untreated children in the same year.*

Parameter	Untreated 2017	Treated 2017	Δ%	P
Rhinitis	1.5±0.9	0.4±0.7	-73.3	0.0121
Watery eyes	0.9±1.1	0.1±0.3	-88.9	0.0358
Cough/bronchospasm	1.5±1.4	0.6±0.9	-60.0	0.11

*Expressed as mean±standard deviation with: 0 = no signs; 1 = mild signs; 2 = moderate signs; 3 = severe signs.

TABLE II.—*Characteristics of children in the treatment arm of the study.*

Group	Sex	N.	Age*	P
Untreated 2016	M	8	10.8±3.5	NS
	F	3	8±1	NS
Treated 2017	M	8	11.8±3.5	NS
	F	3	9±1	NS
Untreated 2107	M	7	9.9±2.9	NS
	F	4	8.3±3.8	NS

NS: not significant.

*Expressed in years (mean±standard deviation).

TABLE IV.—*Signs and symptoms of atopy according to a visual analogue scale* in children in the prophylaxis arm of the study: comparison between children treated in 2017 with the same subjects (untreated) in 2016.*

Parameter	2016	2017	Δ%	P
Rhinitis	2.5±1.3	1.2±1	-52.0	0.0001
Watery eyes	1.9±1.5	0.8±0.9	-57.9	0.0001
Cough/bronchospasm	1.9±1.3	1.1±1.1	-42.1	0.0002

*Expressed as mean±standard deviation with: 0 = no signs; 1 = mild signs; 2 = moderate signs; 3 = severe signs.

same year (2017) were compared, a significant ($P<0.05$) reduction in rhinitis and watery eyes was seen; when the same children from two different years (2017 *versus* 2016) were compared, only cough/bronchospasm was significantly reduced ($P<0.05$).

Analysis of signs, symptoms and drug use in children administered the probiotic mixture (both prophylactically and not) in 2017 compared with untreated children in the same year, showed a significant association between the use of BB12 and L3 and a reduction in rhinitis, watery eyes and cough/bronchospasm ($OR=0.2$) and a decrease in the number of days of treatment with conventional drugs such as oral corticosteroids

TABLE VI.—*Signs and symptoms of atopy according to a visual analogue scale* in children in the treatment arm of the study: comparison between children treated in 2017 and same subjects (untreated) in 2016.*

Parameter	2016	2017	Δ%	P
Rhinitis	1.1±1.3	0.4±0.7	-63.6	0.125
Watery eyes	0.8±1.3	0.1±0.3	-87.5	0.125
Cough/bronchospasm	2.1±1.3	0.6±0.9	-70.0	0.0156

*Expressed as mean±standard deviation with: 0 = no signs; 1 = mild signs; 2 = moderate signs; 3 = severe signs.

TABLE VII.—*Odd ratio (OR) in all children in the study: comparison between different subjects in the same year (untreated 2017 versus treated 2017).*

Parameter	Untreated 2017	Treated 2017	P	OR (95% CI)
Rhinitis	0.9±0.3	0.6±0.5	0.004	0.2 (0.1-0.6)
Watery eyes	0.7±0.4	0.4±0.5	0.002	0.2 (0.1-0.6)
Cough/bronchospasm	0.8±0.4	0.5±0.5	0.005	0.2 (0.1-0.7)
Antihistamine (oral)	0.7±0.5	0.5±0.5	0.09	0.5 (0.2-1.1)
Corticosteroid (oral)	0.1±0.4	0±0.1	0.044	0.5 (0.2-0.8)
Antileukotrienes (oral)	0.1±0.3	0±0.1	0.29	0.3 (0.1-3.0)
Corticosteroid (inhaled)	0.3±0.5	0.3±0.5	0.82	1.1 (0.5-2.7)
Adrenergic β2 agonists	0.3±0.5	0.2±0.4	0.26	0.6 (0.2-1.5)
Antihistamine (eye drops)	0.1±0.3	0.1±0.3	0.85	1.1 (0.3-4.0)

*Expressed as mean±standard deviation.

TABLE VIII.—*Odds ratio (OR) in all children in the study: comparison between the same subjects in different years (untreated 2016 versus treated 2017).*

Parameter	Untreated 2016*	Treated 2017*	P	OR (95% CI)
Rhinitis	0.8±0.4	0.6±0.5	0.12	0.5 (0.2-1.2)
Watery eyes	0.6±0.5	0.4±0.5	0.07	0.5 (0.2-1.0)
Cough/bronchospasm	0.7±0.4	0.5±0.5	0.035	0.4 (0.2-0.9)
Antihistamine (oral)	0.8±0.4	0.5±0.5	0.002	0.2 (0.1-0.6)
Corticosteroid (oral)	0±0.2	0±0.1	0.57	0.5 (0.1-5.6)
Antileukotrienes (oral)	0.1±0.3	0±0.1	0.18	0.2 (0.1-2.2)
Corticosteroid (inhaled)	0.6±0.5	0.3±0.5	0.008	0.3 (0.1-0.7)
Adrenergic β2 agonists	0.4±0.5	0.2±0.4	0.06	0.4 (0.2-0.6)
Antihistamine (eye drops)	0.2±0.4	0.1±0.3	0.41	0.6 (0.2-1.9)

*Expressed as mean±standard deviation.

(OR=0.5) (Table VII). Comparison of the findings in the same group of children untreated in 2016 but treated in 2017, showed a significant association between the use of probiotics and a reduction in cough/bronchospasm (OR=0.4) and a decrease in the number of days of treatment with conventional drugs such as oral antihistamines, inhaled corticosteroids and adrenergic β2 agonists (OR=0.2, OR=0.3 and OR=0.4, respectively) (Table VIII).

Discussion

B. animalis subsp. *lactis* BB12 (BB12) is a well-documented probiotic. It has been described in more than 300 scientific papers including over 130 clinical trial reports. Its complete DNA sequence has been determined and published. It has excellent gastric acid resistance, bile tolerance, gut adherence and colonizing properties in newborn infants, children and adults.¹¹ Immunologically, it seems to promote Th1 response as shown clinically following vaccination against

influenza¹² or in reducing the risk of respiratory tract infection in early childhood.¹³ As known, immune response are polarized either *versus* a Th1-type or *versus* a Th2-type, being this last the typical pro-atopy response.¹⁴ The BB12 strain has indeed anti-Th2 properties as shown in a report demonstrating the strain modified allergic inflammation.¹⁵ BB12 can change the immune response so that it does not cause atopic manifestations, as clearly demonstrated in a neonatal gnotobiotic piglet disease model.¹⁶ *E. faecium* L3 (L3) was originally isolated in the feces of a healthy newborn by Russian researchers who identified it through its ability to kill *Streptococcus agalactiae* by mean of enterocins (A and B) release.¹⁷ Recently, a trial in Italy in pregnant women confirmed its ability to reduce swab test positivity to group B streptococcus while also decreasing the number of cases of premature rupture of membranes.¹⁸ The L3 strain has been extensively used in newborn and preterm infants with positive effects on bodyweight, body mass index, respiratory infections, necrotizing entero-

colitis and infective complications.¹⁹⁻²¹ For these reasons we examined the ability of a mixture of BB12 and L3 to reduce the signs and symptoms of atopy in children: *Bifidobacteria* are frequently depleted in atopic children⁷ and adults,⁸ and L3 promotes the preservation of endogenous gut *Bifidobacteria* in children.¹⁰ The results of our retrospective clinical analysis show that this mixture of strains has beneficial effects in atopic children, especially when administered prophylactically at least 3 months before the appearance of the signs and symptoms of allergy and/or asthma. The fact that the mixture shows better results when administered prophylactically indicates that time is required for the probiotics to colonize the gut and exert their beneficial effects. We compared the results in the treated group with the results in the two control groups: the same children in the previous year (2016) and different children in the same year (2017). As atopy can improve or worsen on an annual basis due to variations in parameters such as weather conditions and pollen quantity, the use of two control groups reduced the risk of incorrect conclusions. As far as we are aware, there was less atopy in 2017 compared with 2016. However, in the prophylactic arm of the study, we did not find any differences in signs and symptoms of rhinitis, watery eyes and cough/bronchospasm between the treated group and the untreated group of the year before (2016) or between the treated and untreated groups of the same year (2017). It is difficult to interpret the findings in the treatment arm of the study because of the small number of children enrolled. Nevertheless, there was a clear tendency for the use of the probiotic mixture to reduce, even if only mildly significantly, the signs and symptoms of atopy. In order to obtain more data, we evaluated the odd ratios of having signs and symptoms of atopy and of using fewer conventional antiatopy drugs by combining all participants from the prophylactic and treatment arms of the study. Better results regarding signs and symptoms were obtained by comparing different children in the same year (treated *versus* untreated in 2017), while better results regarding reduced drug use were obtained by comparing the same children in two different years (treated 2017 *versus* untreated 2016). We do not have a

clear explanation for these differences, but they may be due to an insufficient number of enrolled children, which is a possible source of bias. Another important limitation is the lack of blinding. A prospective rather than a retrospective approach would also have allowed a better explanation of the obtained results.

Although various publications have suggested that gut dysbiosis, often focusing on bifidobacterial species, may underlie atopy and several trials have been conducted where probiotics were administered to atopic subjects, there are few consistent results and meta-analysis does not clearly support to the benefits of probiotic use.²²⁻²⁴ Probiotics seem to be helpful in improving symptoms and quality of life in patients with allergy and/or asthma, but current evidence is strongly limited by study heterogeneity and the use of different outcome measures. Results are often contradictory and there are no guidelines on the use of probiotics to prevent or treat atopy. However, our study suggests that prophylactic administration of the correct probiotics could have a beneficial effect on atopy.

Conclusions

In conclusion, despite possible bias in our retrospective analysis, the prophylactic use in atopic children of a mixture of the widely documented probiotic strains BB12 and L3, reduced signs and symptoms of atopy such as rhinitis, watery eyes and cough/bronchospasm. This reduction resulted in decreased use of conventional drugs including oral antihistamines, inhaled corticosteroids and adrenergic $\beta 2$ agonists.

References

1. Fujimura KE, Lynch SV. Microbiota in allergy and asthma and the emerging relationship with the gut microbiome. *Cell Host Microbe* 2015;17:592–602.
2. Mahowald MA, Rey FE, Seedorf H, Turnbaugh PJ, Fulton RS, Wollam A, *et al.* Characterizing a model human gut microbiota composed of members of its two dominant bacterial phyla. *Proc Natl Acad Sci USA* 2009;106:5859–64.
3. Rutten NB, Gorissen DM, Eck A, Niers LE, Vlieger AM, Besseling-van der Vaart I, *et al.* Long term development of gut microbiota composition in atopic children: impact of probiotics. *PLoS One* 2015;10:e0137681.
4. van Nimwegen FA, Penders J, Stobberingh EE, Postma

DS, Koppelman GH, Kerkhof M, *et al.* Mode and place of delivery, gastrointestinal microbiota, and their influence on asthma and atopy. *J Allergy Clin Immunol* 2011;128:948–55.e1.

5. Abrahamsson TR, Jakobsson HE, Andersson AF, Björkstén B, Engstrand L, Jenmalm MC. Low gut microbiota diversity in early infancy precedes asthma at school age. *Clin Exp Allergy* 2014;44:842–50.

6. Björkstén B, Sepp E, Julge K, Voor T, Mikelsaar M. Allergy development and the intestinal microflora during the first year of life. *J Allergy Clin Immunol* 2001;108:516–20.

7. Fujimura KE, Sitarik AR, Havstad S, Lin DL, Levan S, Fadrosch D, *et al.* Neonatal gut microbiota associates with childhood multisensitized atopy and T cell differentiation. *Nat Med* 2016;22:1187–91.

8. Hevia A, Milani C, López P, Donado CD, Cuervo A, González S, *et al.* Allergic patients with long-term asthma display low levels of *Bifidobacterium adolescentis*. *PLoS One* 2016;11:e0147809.

9. Ruiz L, Delgado S, Ruas-Madiedo P, Sánchez B, Margolles A. *Bifidobacteria* and their molecular communication with the immune system. *Front Microbiol* 2017;8:2345.

10. Lo Skiavo LA, Gonchar NV, Fedorova MS, Suvorov AN. [Dynamics of contamination and persistence of *Clostridium difficile* in intestinal microbiota in newborn infants during antibiotic therapy and use of probiotic strain *enterococcus faecium* L3]. *Antibiot Khimioter* 2013;58:13–8. Russian.

11. Jungersen M, Wind A, Johansen E, Christensen JE, Stuer-Lauridsen B, Eskesen D. The Science behind the Probiotic Strain *Bifidobacterium animalis* subsp. *lactis* BB-12®. *Microorganisms* 2014;2:92–110.

12. Rizzardini G, Eskesen D, Calder PC, Capetti A, Jespersen L, Clerici M. Evaluation of the immune benefits of two probiotic strains *Bifidobacterium animalis* ssp. *lactis*, BB-12® and *Lactobacillus paracasei* ssp. *paracasei*, L. casei 431® in an influenza vaccination model: a randomised, double-blind, placebo-controlled study. *Br J Nutr* 2012;107:876–84.

13. Taipale TJ, Pienihäkkinen K, Isolauri E, Jokela JT, Söderling EM. *Bifidobacterium animalis* subsp. *lactis* BB-12 in reducing the risk of infections in early childhood. *Pediatr Res* 2016;79:65–9.

14. Nagai H. Recent research and developmental strategy of anti-asthma drugs. *Pharmacol Ther* 2012;133:70–8.

15. Isolauri E, Arvola T, Sütas Y, Moilanen E, Salminen S. Probiotics in the management of atopic eczema. *Clin Exp Allergy* 2000;30:1604–10.

16. Chattha KS, Vlasova AN, Kandasamy S, Rajashekara G, Saif LJ. Divergent immunomodulating effects of probiotics on T cell responses to oral attenuated human rotavirus vaccine and virulent human rotavirus infection in a neonatal gnotobiotic piglet disease model. *J Immunol* 2013;191:2446–56.

17. Ermolenko EI, Chernysh AI, Martsinkovskaia IV, Suvorov AN. [Influence of probiotic enterococci on the growth of *Streptococcus agalactiae*]. *Zh Mikrobiol Epidemiol Immunobiol* 2007;(5):73–7. Russian.

18. Di Pierro F, Parolari A, Brundu B, Nigro R. Positive clinical outcomes derived from using a proprietary mixture of selected strains during pregnancy. *Acta Biomed* 2017;87:259–65.

19. Gonchar NV, Alekhina LA, Suvorov AN. [Probiotic strains of enterococci as a means of therapy and prophylaxis of intestinal diseases in children (review of literature)]. *Eksp Klin Gastroenterol* 2013;(1):74–8. Russian.

20. Lo Skiavo LA, Gonchar NV, Suvorov AN, Shabalov NP, Grigor'ev SG. [Value of probiotic in reducing frequency of infectious complications in premature infants]. *Antibiot Khimioter* 2014;59:30–5. Russian.

21. Gonchar NV, Suvorov AN, Maryshev VP, Sorokina TM, Churkova TV, Kharit SM. Probiotics, nutritional status and resistance to respiratory infections in infants. *Eksp Klin Gastroenterol* 2015;(1):48–54. Russian.

22. Elazab N, Mendy A, Gasana J, Vieira ER, Quizon A, Forno E. Probiotic administration in early life, atopy, and asthma: a meta-analysis of clinical trials. *Pediatrics* 2013;132:e666–76.

23. Zajac AE, Adams AS, Turner JH. A systematic review and meta-analysis of probiotics for the treatment of allergic rhinitis. *Int Forum Allergy Rhinol* 2015;5:524–32.

24. Zhang GQ, Hu HJ, Liu CY, Zhang Q, Shakya S, Li ZY. Probiotics for prevention of atopy and food hypersensitivity in early childhood: a PRISMA-compliant systematic review and meta-analysis of randomized controlled trials. *Medicine (Baltimore)* 2016;95:e2562.

Conflicts of interest.—Francesco Di Pierro is on the Scientific Board of Omeopiacenza, the firm supplying the product in Italy.

Article first published online: April 12, 2018. - Manuscript accepted: April 6, 2018. - Manuscript received: January 23, 2018.